

The First Microsatellite-based Characterisation of *Blastocystis* sp. ST3 Isolates and Population Structure Analysis

Blastocystis sp. ST3 İzolatlarının Mikrosatellit Temelli İlk Karakterizasyonu ve Popülasyon Yapısının Analizi

İ Sema Ertuğ¹, İ Erdoğan Malatyali¹, İ Hatice Ertabaklar¹, İ Bülent Bozdoğan², İ Mahmut Sincen³, İ Özgür Güçlü⁴

¹Aydın Adnan Menderes University Faculty of Medicine, Department of Parasitology, Aydın, Türkiye

²Aydın Adnan Menderes University Faculty of Medicine, Department of Microbiology, Aydın, Türkiye

³Aydın Adnan Menderes University Faculty of Engineering, Department of Computer Engineering, Aydın, Türkiye

⁴Aydın Adnan Menderes University, Sultanhisar Vocational School, Department of Plant and Animal Production, Aydın, Türkiye

Cite this article as: Ertuğ S, Malatyali E, Ertabaklar H, Bozdoğan B, Sincen M, Güçlü Ö. The first microsatellite-based characterisation of *Blastocystis* sp. ST3 isolates and population structure analysis. Türkiye Parazitolojisi Dergisi. 2025;49(1):17-22.

ABSTRACT

Objective: *Blastocystis* sp. is an intestinal Stramenopile that can infect both humans and animals. Genetic variability of *Blastocystis* has been investigated with a variety of molecular methods and different subtypes (ST) have been identified. The present study aimed to characterize microsatellite loci and population structure for *Blastocystis* sp. ST3, the most common ST in human faecal samples.

Methods: The genome of *Blastocystis* sp. ST3 in GenBank was analyzed for the presence of microsatellites containing at least eight repeat units. Specific primers were designed for each locus and polymorphisms were identified using bioinformatics tools. The population structure was determined, and microsatellite typing was conducted on 18 *Blastocystis* sp. ST3 genomic DNA samples from the routine laboratory at Aydın Adnan Menderes University Hospital.

Results: The whole-genome scan of *Blastocystis* sp. ST3 revealed 12 microsatellite loci with at least eight repeats. All loci were successfully amplified with the designed primers and eight of them were selected for genotyping. Microsatellite polymorphism analysis showed that each isolate had a unique profile (18 isolates, 18 different microsatellite types). Furthermore, the isolates were grouped into two distinct population clusters.

Conclusion: *Blastocystis* sp. ST3 isolates exhibited significant variability in their microsatellite repeats. The polymorphic microsatellite loci offer a novel approach to study the detailed genetic diversity and population structure of *Blastocystis* sp. ST3.

Keywords: *Blastocystis* sp., genetic diversity, microsatellite, polymorphism

Öz

Amaç: *Blastocystis* sp. hem insanları hem de hayvanları enfekte edebilen intestinal yerleşimli bir Stramenopil türü olarak sınıflandırılmaktadır. Genetik çeşitliliği farklı moleküler yöntemlerle araştırılmış ve çok sayıda alt tipi (ST) tanımlanmıştır. Bu çalışmada insan dışkı örneklerinde en yaygın saptanan *Blastocystis* sp. ST olan ST3'e özgü mikrosatellit lokusların karakterizasyonu ve popülasyon yapısının belirlenmesi amaçlanmıştır.

Yöntemler: *Blastocystis* sp. ST3'ün tüm genom dizisi taranarak en az sekiz tekrarlı mikrosatellit lokusları belirlenmiştir. Her bir lokusa spesifik primerler dizayn edilmiş ve polimorfizmler biyoinformatik araçlar ile analiz edilmiştir. Aydın Adnan Menderes Üniversitesi Hastanesi rutin laboratuvarından elde edilen 18 *Blastocystis* sp. ST3 genomik DNA örneği kullanılarak popülasyon yapısı belirlenmiş ve mikrosatellit tiplendirme gerçekleştirilmiştir.

Bulgular: *Blastocystis* sp. ST3 tüm genom taraması sonucu en az sekiz tekrarlı 12 mikrosatellit lokusu tespit edilmiştir. Dizayn edilen primerler ile lokusların tümü başarıyla amplifiye edilmiş ve bunlardan sekizi genotiplendirmede kullanılmak üzere seçilmiştir. *Blastocystis* sp. ST3 izolatları mikrosatellit polimorfizmlerine göre tiplendirildiğinde her bir izolat farklı bir profil (18 izolat, 18 farklı mikrosatellit tipi) göstermiştir. Ayrıca izolatların iki ayrı popülasyon grubunda toplandığı saptanmıştır.



Received/Geliş Tarihi: 24.11.2024 Accepted/Kabul Tarihi: 15.04.2025 Epub: 18.04.2025 Publication Date/Yayınlanma Tarihi: 09.05.2025

Address for Correspondence/Yazar Adresi: Erdoğan Malatyali, Aydın Adnan Menderes University Faculty of Medicine, Department of Parasitology, Aydın, Türkiye

E-mail/E-Posta: erdogan.malatyali@adu.edu.tr ORCID ID: orcid.org/0000-0002-3943-467X



©Copyright 2025 Turkish Society for Parasitology - Available online at www.turkiyeparazitolog.org

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial (CC BY-NC-ND) 4.0 International License.

Sonuç: *Blastocystis* sp. ST3 izolatlarının mikrosatellit tekrarları açısından birbirlerinden oldukça farklı bir yapıya sahip oldukları belirlenmiştir. Polimorfik mikrosatellit lokusları, *Blastocystis* sp. ST3 izolatlarının detaylı genotiplendirilmesine ve popülasyon yapılarının araştırılmasına olanak sağlayan yeni bir yaklaşım olarak görülmektedir.

Anahtar Kelimeler: *Blastocystis* sp., genetik çeşitlilik, mikrosatellit, polimorfizm

INTRODUCTION

Blastocystis is one of the most common protozoa in the human gastrointestinal system. Most researchers have agreed that *Blastocystis* transmission occurs via the faecal-oral route, primarily through the ingestion of cyst forms in water or food (1). Zoonotic transmission between a variety of animals and humans may also be possible (2). Many subtypes (STs) have been identified to date mostly relying on 18S rRNA coding gene polymorphisms, ST1-4 accounts for the great majority of human samples in many studies (3-5). The genetic diversity of *Blastocystis* is among the most interesting topics in recent years because it has a great contribution to host specificity, pathogenicity, experimental studies and also microbiota (5,6). Amplification of specific parts of rRNA genes and sequencing is the most common method for detecting genotypes. Moreover, Matrix-assisted laser desorption/ionisation mass spectrometry (MALDI MS) has been used for genotype determination (7).

Microsatellites are 2-9 bp length genetic markers that can be found abundantly in the genomes of eukaryotic organisms. They allow the identification of the causative agents precisely in populations of different genotypes because they contain high amounts of genetic diversity (8). Microsatellite markers are generally regarded as neutral alleles, which makes them ideal markers for determining the history of populations. In addition, up to today, MS markers have been used in several parasitic protozoa including *Leishmania tropica*, *Trypanosoma* sp., *Plasmodium* sp., and *Toxoplasma gondii* (9-12).

The development and application of alternative genotyping methods may contribute to the understanding of controversial issues in *Blastocystis* sp. pathogenicity such as pathogenicity. In the present study, we aimed to characterize the microsatellite loci for *Blastocystis* sp. ST3 for the first time in the literature and analyse the genetic structure of *Blastocystis* sp. ST3 isolates.

METHODS

Determination of Microsatellite Markers and Primer Design

The complete genomic sequence of a human *Blastocystis* sp. ST3 isolate, an assemblage of 917 partial sequences, was acquired from the National Center for Biotechnology Information database (Genbank, Acc. no: JZRK00000000). Microsatellites mining was carried out using the software Msatcommander 0.8.1 and http://insilico.ehu.es/mini_tools/microsatellites/website with the motif criteria, di-, tri-, tetra-, penta-, and hexanucleotide repeats of microsatellites at least 8 within the complete genome of *Blastocystis* sp. ST3 (13). Primers for microsatellite loci (n=2) were designed with Primer3, the online version, with default settings (14,15). For potential repetitive elements other than perfect repeats, the flanking regions were also analysed. In order to identify robust loci with an annealing temperature of at least 55 °C, approximately 150-200 bp flanking each side of the repeat were included.

Blastocystis sp. ST3 Isolates

The ethical approval from a Local Ethical Committee in Aydın Adnan Menderes University Faculty of Medicine (no: 2015/10) was obtained. A total of 18 *Blastocystis* sp. ST3 isolates were used in the study. The isolates were previously acquired by culturing direct microscopy-positive faecal samples in the routine diagnostic laboratory at Aydın Adnan Menderes University Hospital. The cultures in 3 mL of Jones medium were subjected to genomic DNA isolation with DNAzol kit (Invitrogen, Life Technologies). *Blastocystis* sp. SSU-rDNA gene was partially amplified with the primers RD5 and BhRD_r in a single round of polymerase chain reaction (PCR) (16). The amplicons were sequenced, and ST were detected using the *Blastocystis* sp. sequence typing database (pubmlst.org/*Blastocystis*) (17).

PCR Amplification and Genotyping

The optimal annealing temperature for the PCR amplification for each microsatellite locus was determined with gradient PCR. The fluorophores FAM and HEX were used to label the 5' ends of the forward primer. The amplifications were performed in 30 µL of volume: 0.5 mM each of the primers, 1.5 mM MgCl₂, 2.5 mM dNTP, 1.0 U Taq polymerase, and 1-2 µL of template DNA. The reaction for each locus was set as follows: 95 °C for 5 min, 35 cycles (at 95 °C for 30 s, 55-60 °C for 30 s and 72 °C for 45 s), and a final extension at 72 °C for 7 min. The length of alleles was detected with an automatic sequencer.

Statistical Analysis

The allele sizes of microsatellites were exactly determined with Genemarker 2.6.3 (Soft Genetics LLC, USA). The calculation of genetic variation in microsatellite loci was performed with GenAlEx 6.5 (18). GENEPOP 3.3 software was used to detect genotypic linkage disequilibrium between pairwise loci (19). GenAlEx 6.5 program was used to detect the allele numbers (N_A), effective allele numbers (N_E), the frequencies of alleles, intra-population diversity of alleles, and pairwise comparisons of the isolates (18). The expected heterozygosity (H_E) of loci was calculated using the Arlequin 3.11 (20). In multiple loci, haplotype overlaps were determined with GenAlEx 6.5 (18).

Population Structure and Microsatellite Typing

Population structure was analysed with a Bayesian clustering method in STRUCTURE ver. 2.3 program (21). The admixture model with correlated allele frequency parameters was used to detect the estimated number of genetic clusters (K). Ten runs were performed for each K value (ranging from 1 to 10) with 100.000 MCMC repetitions and a burn-in periods of 10,000. The ad hoc estimated likelihood of K (ΔK) was included in the determination of the most likely number of populations (K) based on the rate of change in the log probability of the data [$\ln Pr(X/K)$] (22). Structure Harvester version 0.6.94 was used to infer the most likely number of genetic clusters (K) with both the Evanno and Delta K methods (23). The first isolate was defined as MT1 and

the remaining different isolates (at least one different allele type) were annotated with a new MT number.

RESULTS

Microsatellite Variability and Population Structure

We detected 12 microsatellite loci with at least 8 repeats in *Blastocystis* sp. ST3 genome and of these 11 loci were polymorphic. All loci have three repeats, and most were characterized with (GAT)_n and (ATC)_n (Table 1).

There was no significant linkage disequilibrium when the loci were compared dually ($p > 0.05$). The total number of alleles (N_A) for each locus changed from three to 11, the average was 6.55 alleles per locus (Table 1). The mean effective number of alleles (N_E) was 4.06 (range from 1.841 to 8.526). BHST3-142 loci ($h = 0.457$) provided the minimum intra-population diversity, while BHST3-838 locus ($h = 0.883$) provided the highest. The average expected heterozygosity value was determined as 0.740 ± 0.146 (min: 0.399 and max: 0.935). Two distinct genetic groups were suggested with the analyses of microsatellite data by a structure clustering algorithm ($K = 2.297$) (Figure 1).

Microsatellite Typing

We selected eight loci: BH142, BH217, BH278, BH302, BH399, BH476, BH806 and BH808 for MT typing of *Blastocystis* sp. ST3 isolates. These loci were selected relying on the reliability of microsatellite length, clarity in representations, and amplification/reproducibility. Microsatellite based typing of isolates revealed that each isolate had a unique MT type (Table 2).

DISCUSSION

Blastocystis sp. is a common enteric stramenophile with a high genetic diversity between isolates and an unclear role in the development of intestinal diseases and pathogenicity. The molecular typing of *Blastocystis* sp. isolates is valuable to study its population structure of parasites on a global scale. Genetically diverse isolates may influence the frequency of *Blastocystis* sp., for example, host specificity and adaptation to novel hosts (24). In addition, several studies reported a decreased bacterial diversity and changed intestinal microbiome in certain *Blastocystis* sp. ST (6). The development of novel genotyping methods is important for an improved understanding of the genotype distribution of *Blastocystis* sp. isolates and phenotypic characteristics. In the

Table 1. Characterization of 12 microsatellite markers for the *Blastocystis* sp. ST3

Locus	Primer sequence (5'-3')	Size	N_A	Motif	N_E	h	H_E	GenBank accession number
BHST3-142	F: TACAGCTGTTCCACCCACTC R: CTCCCCTCTCCCCTAGTCAC	216-234	4	(GAT) ¹³	1.841	0.457	0.399	JZRK01000142
BHST3-217	F: GGAGGGAGGTTCTTGGTCTC R: AGAGGGATGTGGTGGAGTTG	232-349	6	(ATC) ¹⁰	3.767	0.735	0.778	JZRK01000217
BHST3-278	F: CCTCCTTGCGCTTCTGAC R: AGGAGGCGGAAAGGAGAAAG	385-409	6	(AGC) ¹¹	3.375	0.704	0.745	JZRK01000278
BHST3-302	F: GAGATGCGACGATTCTCCTC R: GATTTCGCGTTCAATGGTTCT	298-325	6	(ATC) ¹¹	3.600	0.722	0.765	JZRK01000302
BHST3-366	F: ATGGTCAATTGGCTGATGGTG R: AGTTTCTGAGCCACATGGAGA	141-201	9	(ATC) ¹²	4.154	0.759	0.804	JZRK01000366
BHST3-399	F: ATTCTTCGCACGATTCTTCG R: TCATGGAGGTGGTCAACAAA	364-370	3	(GAT) ⁸	2.160	0.537	0.569	JZRK01000399
BHST3-476	F: AGAAGAAGCTCTTCGCGTTG R: CAGATCCGCTAGGTCATTT	321-357	7	(AAG) ⁸	4.050	0.753	0.811	JZRK01000476
BHST3-500	F: GTGGTTGAGGAGGAGATGA R: GAGAGGAGGCGTCGATGATA	-	-	(GAT) ⁸				JZRK01000500
BHST3-654	F: ATTGGTGATCGTGTGGTGA R: TGGTGACGAGTTCGATGAAG	297-471	11	(ATC) ¹¹	6.480	0.846	0.895	JZRK01000654
BHST3-806	F: GACGTGGGTGAGGAGGATTA R: GTTGAGAGTTCGGGGGTAT	367-385	6	(GAT) ⁹	4.263	0.765	0.811	JZRK01000806
BHST3-808	F: TGACAGTTGCTCCTTACCC R: TTGGAGTCGTTGGACATTGA	304-325	4	(ATC) ⁹	2.492	0.599	0.634	JZRK01000808
BHST3-838	F: TAAATGTCGGAGGGAAGGTG R: GGGAGATGAGTGCATTGACA	158-295	10	(GAT)8(GTT) ⁸	8.526	0.883	0.935	JZRK01000838
Mean			6.55		4.064	0.705	0.747	

N_A : Observed number of alleles, I : Shannon's information index, N_E : Effective number of alleles; H_E : Expected heterozygosity




Figure 1. A) Bayesian clustering analysis (K=2) based on SSR data for the studied *Blastocystis* sp. ST3 (18 isolates), vertical black lines separate isolates, **B)** Scatter plot of the possible cluster numbers (K; horizontal axis) against the ad-hoc estimated likelihood of ΔK (vertical axis) by changing the likelihood rate (circle)

present study, we characterised the microsatellite markers for the most common *Blastocystis* sp. subtype in human faecal samples, namely ST3. The most significant finding from this study was that the microsatellite loci in *Blastocystis* sp. ST3 isolates were highly variable, with each of the 18 isolates exhibiting a distinct or unique microsatellite type. A great genetic difference between *Blastocystis* sp. ST was noted in the literature (25). The comparison of *Blastocystis* sp. ST1, ST4 and ST7 genomes found great divergence in terms of genome assembly size, number of protein-coding genes, guanine-cytosine content, average gene size (bp) and also number of introns (25,26). In addition to subtype level genetic difference in *Blastocystis* sp., the isolates were highly separated in terms of intra-subtype variation. A previous study investigated the intra-subtype genetic diversity and host specificity of *Blastocystis* sp. ST3 with multilocus sequence typing including SSU-rDNA and mitochondrion-like organelle sequences (27). The authors reported that human isolates were restricted to a single clade suggesting relatively high host specificity of ST3 human-to-human transmission. Another study reported that 11 human *Blastocystis* sp. ST3 isolates were distributed in nine sequence types with the same method. There were only three isolates that shared the same sequence type (28). However, the use of a single gene, most commonly SSU rRNA, detected a limited intra-subtype genetic diversity in *Blastocystis* sp. ST3 STs. For example, intra-subtype genetic polymorphisms were identified five in ST3 isolates in China and the similarity of SSU rRNA sequences was 99.8% in Iran (29,30).

In the present study, the population structure analysis of *Blastocystis* sp. ST3 isolates revealed two distinct groups. The genomic DNA samples in our study were limited to a specific location, the same hospital laboratory. Collecting samples

Table 2. The microsatellite types (MT) of *Blastocystis* sp. ST3 isolates

Isolate	Microsatellite loci								MT
	142	217	278	302	399	476	806	808	
1	3	5	2	3	2	5	6	3	MT1
2	3	2	4	5	1	1	4	3	MT2
3	3	1	6	1	1	5	4	2	MT3
4	3	2	5	2	1	5	6	2	MT4
5	5	6	4	2	1	1	6	3	MT5
6	3	3	3	6	2	1	5	4	MT6
7	4	6	3	1	2	6	3	3	MT7
8	3	3	3	3	2	4	1	2	MT8
9	3	4	5	3	2	3	2	2	MT9
10	1	2	3	3	1	5	6	1	MT10
11	3	2	5	3	1	2	2	2	MT11
12	3	5	3	3	2	3	2	2	MT12
13	3	2	3	3	2	1	6	2	MT13
14	3	2	5	4	2	5	4	3	MT14
15	1	2	3	2	3	2	5	3	MT15
16	2	4	1	3	2	5	6	2	MT16
17	3	5	3	2	2	2	3	3	MT17
18	3	2	5	5	2	5	6	3	MT18

from various localities, such as different cities or countries, will enable a detailed phylogenetic analysis of microsatellites in *Blastocystis* sp. ST3. The major contribution of the present study is the characterization of microsatellites in *Blastocystis* sp. ST3 and the presentation of the associated methodology. This characterization has important implications for future research and practices, particularly in areas such as zoonotic transmission, microbiota interactions, and pathogenesis. These are the widely discussed topics in the literature on *Blastocystis* (5). Microsatellite polymorphisms have previously been described as valuable genetic markers for investigating zoonotic transmission in both parasitic microorganisms. For instance, certain microsatellite alleles in *Cryptosporidium parvum* have been associated with human-adapted or zoonotic strains (31). Similarly, microsatellite markers have been used to compare *Plasmodium falciparum* isolates from patients with cerebral malaria and those with uncomplicated malaria, to determine if there is a possible link between genetic variation and pathogenicity (32).

CONCLUSION

In conclusion, the characterisation of microsatellite loci in *Blastocystis* sp. ST3 revealed different profiles or types showing high intra-subtype diversity in terms of microsatellite repeats. This novel genotyping approach may be used in molecular epidemiology and population structure screening studies.

*Ethics

Ethics Committee Approval: The ethical approval from a Local Ethical Committee in Aydın Adnan Menderes University Faculty of Medicine (no: 2015/10) was obtained.

Informed Consent: All participants were informed about the study, and their consents were obtained.

Acknowledgments

The study was presented at 3. International *Blastocystis* Conference, 2-4 June 2024, online.

Footnotes

*Authorship Contributions

Concept: S.E., H.E., B.B., Ö.G., Design: S.E., H.E., B.B., Ö.G., Data Collection or Processing: E.M., M.S., Analysis or Interpretation: S.E., E.M., M.S., Ö.G., Literature Search: S.E., E.M., H.E., Writing: S.E., E.M., H.E., Ö.G.

Conflict of Interest: No conflict of interest was declared by the authors.

Financial Disclosure: This study was supported by the Scientific and Technical Research Council of Türkiye (TUBITAK-215S655).

REFERENCES

- Shams M, Shamsi L, Yousefi A, Sadrebazzaz A, Asghari A, Mohammadi-Ghalehbin B, et al. Current global status, subtype distribution and zoonotic significance of *Blastocystis* in dogs and cats: a systematic review and meta-analysis. *Parasit Vectors*. 2022; 15: 225.
- Chen S, Meng W, Shi X, Chai Y, Zhou Z, Liu H, et al. Occurrence, genetic diversity and zoonotic potential of *Blastocystis* sp. in forest musk deer (*Moschus berezovskii*) in Southwest China. *Parasite*. 2022; 29: 34.
- Malatyali E, Ertabaklar H, Ertug S. Subtype distribution of *Blastocystis* in Türkiye. *Turkiye Parazit Derg*. 2023; 47: 184-9.
- Santin M, Figueiredo A, Molokin A, George NS, Koster PC, Dashti A, et al. Division of *Blastocystis* ST10 into three new subtypes: ST42-ST44. *J Eukaryot Microbiol*. 2024; 71: e12998.
- Wawrzyniak I, Poirier P, Viscogliosi E, Dionigia M, Texier C, Delbac F, et al. *Blastocystis*, an unrecognized parasite: an overview of pathogenesis and diagnosis. *Ther Adv Infect Dis*. 2013; 1: 167-78.
- Deng L, Tan KSW. Interactions between *Blastocystis* subtype ST4 and gut microbiota in vitro. *Parasit Vectors*. 2022; 15: 80.
- Martiny D, Bart A, Vandenberg O, Verhaar N, Wentink-Bonnema E, Moens C, et al. Subtype determination of *Blastocystis* isolates by matrix-assisted laser desorption/ionisation time-of-flight mass spectrometry (MALDI-TOF MS). *Eur J Clin Microbiol Infect Dis*. 2014; 33: 529-36.
- Havrylyuk T, Ferreira MU. A closer look at multiple-clone *Plasmodium vivax* infections: detection methods, prevalence and consequences. *Mem Inst Oswaldo Cruz*. 2009; 104: 67-73.
- Ajzenberg D, Banuls AL, Tibayrenc M, Darde ML. Microsatellite analysis of *Toxoplasma gondii* shows considerable polymorphism structured into two main clonal groups. *Int J Parasitol*. 2002; 32: 27-38.
- Schwenkenbecher JM, Wirth T, Schnur LF, Jaffe CL, Schallig H, Al-Jawabreh A, et al. Microsatellite analysis reveals genetic structure of *Leishmania tropica*. *Int J Parasitol*. 2006; 36: 237-46.
- Simo G, Njiokou F, Tume C, Lueong S, De Meeus T, Cuny G, et al. Population genetic structure of Central African *Trypanosoma brucei gambiense* isolates using microsatellite DNA markers. *Infect Genet Evol*. 2010; 10: 68-76.
- Imwong M, Nair S, Pukrittayakamee S, Sudimack D, Williams JT, Mayxay M, et al. Contrasting genetic structure in *Plasmodium vivax* populations from Asia and South America. *Int J Parasitol*. 2007; 37: 1013-22.
- Faircloth BC. msatcommander: detection of microsatellite repeat arrays and automated, locus-specific primer design. *Mol Ecol Resour*. 2008; 8: 92-4.
- Koressaar T, Remm M. Enhancements and modifications of primer design program Primer3. *Bioinformatics*. 2007; 23: 1289-91.
- Untergasser A, Cutcutache I, Koressaar T, Ye J, Faircloth BC, Remm M, et al. Primer3-new capabilities and interfaces. *Nucleic Acids Res*. 2012; 40: e115.
- Scicluna SM, Tawari B, Clark CG. DNA barcoding of *Blastocystis*. *Protist*. 2006; 157: 77-85.
- Jolley KA, Maiden MC. BIGSdb: scalable analysis of bacterial genome variation at the population level. *BMC Bioinformatics*. 2010; 11: 595.
- Peakall R, Smouse PE. GenAlEx 6.5: genetic analysis in Excel. Population genetic software for teaching and research—an update. *Bioinformatics*. 2012; 28: 2537-9.
- Raymond M, Rousset F. An exact test for population differentiation. *Evolution*. 1995; 49: 1280-3.
- Excoffier L, Estoup A, Cornuet JM. Bayesian analysis of an admixture model with mutations and arbitrarily linked markers. *Genetics*. 2005; 169: 1727-38.
- Pritchard JK, Stephens M, Donnelly P. Inference of population structure using multilocus genotype data. *Genetics*. 2000; 155: 945-59.
- Evanno G, Regnaut S, Goudet J. Detecting the number of clusters of individuals using the software STRUCTURE: a simulation study. *Mol Ecol*. 2005; 14: 2611-20.
- Earl DA, vonHoldt BM. STRUCTURE HARVESTER: a website and program for visualizing STRUCTURE output and implementing the Evanno method. *Cons Gen Res*. 2012; 4: 359-61.
- Noel C, Dufernez F, Gerbod D, Edgcomb VP, Delgado-Viscogliosi P, Ho LC, et al. Molecular phylogenies of *Blastocystis* isolates from different hosts: implications for genetic diversity, identification of species, and zoonosis. *J Clin Microbiol*. 2005; 43: 348-55. 3
- Higuera A, Salas-Leiva DE, Curtis B, Patino LH, Zhao D, Jernstrom-Hultqvist J, et al. Draft genomes of *Blastocystis* subtypes from human samples of Colombia. *Parasit Vectors*. 2023; 16: 52.

26. Gentekaki E, Curtis BA, Stairs CW, Klimes V, Elias M, Salas-Leiva DE, et al. Extreme genome diversity in the hyper-prevalent parasitic eukaryote *Blastocystis*. *PLoS Biol.* 2017; 15: e2003769.
27. Stensvold CR, Alfellani M, Clark CG. Levels of genetic diversity vary dramatically between *Blastocystis* subtypes. *Infect Genet Evol.* 2012; 12: 263-73.
28. Ertug S, Malatyali E, Ertabaklar H, Bozdogan B. Multilocus sequence typing of *Blastocystis* isolates in Aydin, Turkey. *Parasitol Int.* 2016; 65: 760-2.
29. Zhao W, Ren G, Wang L, Xie L, Wang J, Mao J, et al. Molecular prevalence and subtype distribution of *Blastocystis* spp. among children who have diarrhea or are asymptomatic in Wenzhou, Zhejiang Province, China. *Parasite.* 2024; 31: 12.
30. Alinaghizade A, Mirjalali H, Mohebbi M, Stensvold CR, Rezaeian M. Inter- and intra-subtype variation of *Blastocystis* subtypes isolated from diarrheic and non-diarrheic patients in Iran. *Infect Genet Evol.* 2017; 50: 77-82.
31. Hunter PR, Hadfield SJ, Wilkinson D, Lake IR, Harrison FC, Chalmers RM. Subtypes of *Cryptosporidium parvum* in humans and disease risk. *Emerg Infect Dis.* 2007; 13: 82-8.
32. Ferreira MU, Nair S, Hyunh TV, Kawamoto F, Anderson TJ. Microsatellite characterization of *Plasmodium falciparum* from cerebral and uncomplicated malaria patients in southern Vietnam. *J Clin Microbiol.* 2002; 40: 1854-7.